


IMAGE

Extinction of auditory threat memory triggers activation of p70 S6 kinase 1 in the basolateral nucleus of the amygdala

TN Huynh^{1,2,3}, E Santini^{1,6}, E Mojica¹, AE Fink¹, BS Hall^{2,3}, RN Fetcho^{2,3}, L Grosenick^{4,5}, K Deisseroth^{4,5}, JE LeDoux¹, C Liston^{2,3} and E Klann¹

Molecular Psychiatry (2018) **23**, 1393; doi:10.1038/mp.2018.8

Phosphorylation of the p70 S6 kinase 1 (S6K1) substrate ribosomal protein S6 at multiple sites is increased following extinction training relative to box controls. The extinction-induced increased S6 phosphorylation is blocked by the S6K1 inhibitor PF-4708671. Please refer to the article by Huynh et al. on pages 1394–1401.

¹Center for Neural Science, New York University, New York, NY, USA; ²Brain and Mind Research Institute, Weill Cornell Medical College, New York, NY, USA; ³Department of Psychiatry, Weill Cornell Medical College, New York, NY, USA; ⁴Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, CA, USA and ⁵Department of Bioengineering, Stanford University, Stanford, CA, USA. Correspondence: Dr E Klann, Center for Neural Science, New York University, 4 Washington Place, Room 809, New York, NY 10003, USA.

E-mail: eklann@cns.nyu.edu

⁶Current address: Department of Neurology, Columbia University, New York, NY 10032, USA.